

Original article

DOI: 10.2478/10004-1254-65-2014-2408

Abnormal sperm morphology in mouse germ cells after short-term exposures to acetamiprid, propineb, and their mixture

Pinar Göç Rasgele

Duzce University, Faculty of Agriculture and Natural Sciences, Department of Biosystems Engineering, Duzce, Turkey

Received in June 2013
CrossChecked in August 2013
Accepted in November 2013

Pesticides are one of the most potent environmental contaminants, which accumulate in biotic and abiotic components of ecosystems. Acetamiprid (Acm), a neonicotinoid insecticide, and Propineb (Pro), a dithiocarbamate fungicide, are widely used to control sucking insects and fungal infections on crops, respectively. The present study was undertaken to investigate the genotoxic effects of these compounds, individually and in mixtures, in mouse germ cells by using the sperm morphology assay. Mice were injected intraperitoneally with 0.625, 1.25, and 2.50 $\mu\text{g mL}^{-1}$ of Acm, 12.5, 25, and 50 $\mu\text{g mL}^{-1}$ of Pro, and their mixture at the same concentrations over 24 and 48 h. Acm did not significantly increase the percentage of abnormal sperm at any concentration. The frequency of abnormal sperm significantly increased after 24 and 48 h of exposure to 50 $\mu\text{g mL}^{-1}$ of Pro. The mixtures of 2.50 $\mu\text{g mL}^{-1}$ of Acm and 50 $\mu\text{g mL}^{-1}$ of Pro induced sperm abnormalities antagonistically both after 24 and 48 h of exposure. Results suggest that Acm was non-genotoxic for mouse germ cells, while Pro may have been a germ cell mutagen due to the observed increase in the frequency of sperm abnormalities. However, to gain better insight into the mutagenicity and DNA damaging potential of both of these pesticides, further studies at molecular level should be done.

KEY WORDS: *combined effects; pesticide mixture; sperm morphology assay*

At present times, the consumption and types of pesticides used in agricultural production is increasing considerably throughout the world due to the rise in population and crop production. As much as 4.6 million tons of pesticides are released annually into the environment and approx. 500 types of these pesticides are considered a threat to the environment and human beings (1).

Pesticides are commonly used in agricultural areas as single compounds or in mixtures. However, many are non-biodegradable and therefore inevitably expose people occupationally or environmentally to pesticide residues, through air, water, and food. Therefore, studies on the mutagenic activity of pesticide mixtures

using different assays on the genetic system of a living organism are vital.

Many investigations have focused on the evaluation of genotoxicity of pesticides in occupational or environmental settings using different methods (2-12). It has been reported that exposure to environmental contaminants including pesticides causes major pathological effects in the human male reproductive system as well as that of experimental animals (13, 14). The sperm morphology assay is one of the most widely used genetic toxicology assays. Furthermore, the development of sperm head abnormality has been used as a reliable short-term biological indicator in the evaluation of chemical genotoxicity (15-17). The

mouse sperm morphology test also has potential in identifying chemicals that induce spermatogenic dysfunction and perhaps heritable mutations (18).

Acetamiprid (Acm), a neonicotinoid insecticide, is widely used to control sucking insects on crops (19). Neonicotinoids are crucially potent neurotoxic insecticides that act as agonists of nicotinic acetylcholine receptors (nAChR) (20). Propineb (Pro), a dithiocarbamate fungicide, is commonly used for disease control in a wide range of crops (21). Dithiocarbamate fungicides are mainly used for the eradication of fungal infections on plants, fruit, and vegetables (22).

Commercial formulations of Acm and Pro are commonly used on agricultural crops such as tomato, potato, melon, apple, and tobacco, either separately or in mixture (23). Mixtures of chemicals are often well-known to be considerably more toxic than its individual components and their combined effect can be additive, synergistic, or antagonistic (9).

Many researchers have reported that pesticides caused genotoxic effects when used individually (8, 10, 11, 15). However, in reality all living organisms are exposed to not only to individual chemicals, but also to their mixtures. Additive, synergistic or antagonistic effects of pesticide mixtures are due to the interactions between the chemicals that comprise them. Thus, in mutagenicity and carcinogenicity studies, the combined use of substances investigated provides results that are much more significant for establishing genotoxic effects. Although there are a few studies focused on the genotoxicity of Acm and Pro (9, 21, 24, 25), the mutagenicity of their mixtures *in vivo* in mouse germ cells of mice has not been investigated, which motivated us to use the sperm morphology assay to perform precisely this type of study.

MATERIALS AND METHODS

Chemicals

In the present study, commercial formulations of Acm (Mosetam 20 SP[®], containing 20 % acetamiprid as active ingredient; CAS No. 135410-20-7) and Pro (Antracol 70 WP[®], containing 70 % propineb as active ingredient; CAS No. 12071-83-9) were used as the test materials. Mosetam 20 SP[®] was obtained from Safa Agriculture (Turkey) and Antracol 70 WP[®] from

Bayer (Turkey). The chemical structures of Acm and Pro are shown in Figures 1 and 2. Giemsa dye (CAS No. 51811-82-6) was obtained from Merck[®] (Darmstadt, Germany). Mitomycin C (MMC; CAS No. 50-07-7) was obtained from Sigma[®] (Taufkirchen, Germany). All test solutions were freshly prepared prior to each experiment.




Figure 1 Chemical structure of acetamiprid




Figure 2 Chemical structure of propineb

Experimental animals

In this study, 90 male mice (*Mus musculus*, 8-10 weeks of age, with average body weight of 20-25 g), were used. They were purchased from Abant Izzet Baysal University Experimental Animals Applications and Research Center, Turkey. The animals were maintained in a closely inbred colony under conventional laboratory conditions at a room temperature of 25±5 °C and in 12 h dark and 12 h light cycles. Food pellets and water were provided *ad libitum*. The experiment was approved by the Ethics Committee of Abant Izzet Baysal University in Turkey.

Study design

The used concentrations of pesticides were selected according to the results of a preliminary study. Based on the pesticide concentrations used to treat plant disease (23). Acm and Pro were dissolved in water in order to obtain the following mixtures: 0.625 µg mL⁻¹ of Acm + 12.5 µg mL⁻¹ of Pro; 1.25 µg mL⁻¹ of Acm + 25 µg mL⁻¹ of Pro; 2.5 µg mL⁻¹ of Acm + 50 µg mL⁻¹ of Pro; 5 µg mL⁻¹ of Acm + 100 µg mL⁻¹ of Pro; 10 µg mL⁻¹ of Acm + 200 µg mL⁻¹ of Pro. It was observed that the mixtures of Acm and

Pro exhibited high cytotoxic effects at two of the highest concentrations (5 µg mL⁻¹ of Acm + 100 µg mL⁻¹ of Pro; 10 µg mL⁻¹ of Acm + 200 µg mL⁻¹ of Pro) and also decreased the ratio of dividing cells at these same concentrations over a 48 h treatment period. Therefore, these two mixtures were discarded and the remaining three were selected to be tested in this study. In addition, Acm and Pro were also tested separately in order to determine whether these pesticides are spermatotoxic individually.

In the present study, mice were intraperitoneally (*i.p.*) injected with Acm and Pro at concentrations of (0.625, 1.25, and 2.50) µg mL⁻¹ and (12.5, 25, and 50) µg mL⁻¹, respectively, over 24 and 48 h. In addition, their mixture was administered at the same test concentrations over the same period.

MMC was used as a positive control, as it was previously confirmed as a mutagenic agent (15). The significant increase in the number of sperm abnormalities in this group showed that the method was correctly applied. Distilled water was used as a negative control. The *i.p.* route was favoured since it is one of the fastest and most efficient means of delivering test chemicals in a short-term assay (26). Ninety male mice were randomly allocated into fifteen groups ($n=3$ per group) and treated over 24 and 48 h. These groups are given in Table 1.

Sperm morphology assay

After the treatment, the animals were sacrificed by cervical dislocation. Both of the cauda epididymises were dissected out, cut into pieces in 5 mL of saline, filtered, and smears were made according to the standard protocol for sperm morphology assay, as proposed by Wyrobek (18, 27). The smears were fixed in methanol and stained with 10 % Giemsa in Sørensen buffer for 10 min. A total of 1000 sperms per animal were scored under a microscope (Olympus CX21®, Japan) with 100x10 magnification. Sperm head abnormalities were determined as having either normal or abnormal morphology according to Wyrobek and Bruce (27). According to these criteria, a “hookless head” does not have a spherical spot at the tip of the sperm head; a “banana head” has a banana-like form; an “amorphous head” lacks the usual hook and is deformed; and a “folded sperm” is folded on itself.

Statistical analysis

The data were analysed by using SPSS 20 for Windows (SPSS Inc., Chicago, IL, USA) and results

Table 1 Study design

Test group	Chemical	Concentration (µg mL ⁻¹)
I	Distilled water	-
II	MMC	0.200
III	Acm	0.625
IV	Acm	1.250
V	Acm	2.500
VI	Distilled water	-
VII	MMC	0.200
VIII	Pro	12.50
IX	Pro	25.00
X	Pro	50.00
XI	Distilled water	-
XII	MMC	0.200
XIII	Acm + Pro	0.625 + 12.50
XIV	Acm + Pro	1.250 + 25.00
XV	Acm + Pro	2.500 + 50.00

All of the treatments were performed over 24 and 48 h

obtained were expressed as mean±standard error (SE) of three individual analyses. The Kruskal-Wallis test was carried out followed by the Mann-Whitney U test to compare the statistical significance of the differences between the treated and control groups. The dose-response relationship was determined using Pearson correlation analysis. $P<0.05$ was considered significant. The measured values were also compared with the expected values. The expected mean value and SE were calculated as follows (28): mean % (expected for Acm + Pro) = mean % (Acm) + mean % (Pro) = 100 % (control); SE (expected for Acm + Pro) = [(SE for Acm)² + (SE for Pro)²]^{1/2}.

Non-parametric Mann-Whitney U test was used to detect the significance of difference between the expected and measured values. Additive, synergistic and antagonistic effects were evaluated to interpret effects of mixtures. When the measured values were insignificantly higher than the expected values, there was an additive effect. When the measured values were significantly higher than the expected values, there was a synergistic effect, whereas when the measured values were significantly lower than the expected ones, there was an antagonistic effect (28).

RESULTS

The results obtained by the sperm morphology assay are presented in Table 2. Acm induced different types of sperm abnormalities such as “hookless”, “banana”, “amorphous”, and “folded” sperms at all concentrations for 24 and 48 h when compared with the negative control (Figure 3), but these increases were not statistically significant. Pro also increased the frequency of abnormal sperm during the same treatment periods. The frequency of abnormal sperm induced by Pro was significant at the highest concentration ($50 \mu\text{g mL}^{-1}$) for 24 and 48 h when compared with the negative controls. As shown in Figure 4, these increases were concentration-dependent (for 24 h: $R^2=0.8033$, $p<0.01$ and for 48 h: $R^2=0.9488$, $p<0.05$). “Hookless head” was the most common abnormality observed at the highest concentration for 24 and 48 h.

It was also observed that the mixture of Acm and Pro did not induce sperm head abnormalities at any of the concentrations over 24 and 48 h when compared with the negative controls. In addition, there was no significant difference between the two time points




Figure 3 a) Normal sperm; b) Without hook (at $50 \mu\text{g mL}^{-1}$ of Pro for 24 h); c) Amorphous head sperm (at the highest concentration of mixture for 48 h) (x1000)

(Acm 24 h vs. Acm 48 h, Pro 24 h vs. Pro 48 h, etc.) regarding abnormal sperm frequency.

The results of individually applied pesticides were also compared to results for their mixture at the same concentrations. Interestingly, it was found that the highest concentration ($2.5 \mu\text{g mL}^{-1}$ of Acm + $50 \mu\text{g mL}^{-1}$ of Pro) significantly decreased the percentage of abnormal sperm for 24 and 48 h when compared with the individual concentrations of Pro ($50 \mu\text{g mL}^{-1}$) in both treatment periods.

In order to detect the possible combined actions of the pesticides, the expected mean value and SE were calculated and measured values were compared to the expected values. According to the results, the mixture of Acm and Pro at treatment durations




Figure 4 Graphs showing concentration-response curve slopes and correlation coefficients (R^2) of % abnormal sperm in mice germ cells treated with Acm, Pro, and their mixture for 24 and 48 h

produced mostly additive effects. However, it was found that the measured abnormal sperm (%) was significantly above the expected values, that is, a synergistic effect was observed at the lowest concentration of the mixture ($0.625 \mu\text{g mL}^{-1}$ of Acm + $12.5 \mu\text{g mL}^{-1}$ of Pro). Also, an antagonistic effect was observed at the highest mixture concentration (Figure 5).

Table 2 Effects of acetamiprid, propineb, and their mixture on sperm morphology in *Mus musculus*

Test group	Abnormal sperm (%)	
	24 h	48 h
I	5.10±0.30	5.13±0.37
II	26.63±0.63*	28.60±0.66*
III	4.00±0.80	3.33±0.69
IV	4.60±0.30	5.06±0.40
V	5.86±0.40	6.40±1.30
VI	4.86±0.35	5.20±0.30
VII	26.70±0.60*	28.60±0.69*
VIII	5.60±0.50	3.33±0.48
IX	6.00±0.50	5.46±0.29
X	11.20±0.91*	10.46±1.23*
XI	4.76±0.39	5.33±0.37
XII	26.40±0.50*	28.80±1.15*
XIII	6.93±2.68	9.66±2.36
XIV	7.13±1.27	3.93±0.69
XV	5.86±1.00 ^a	4.46±0.33 ^a

Data are reported as mean value ± SE

The treatments are explained in detail in Table 1

* p<0.001, when compared with negative controls

^a (p<0.01) each pesticide alone as compared to the mixture of Acm and Pro

DISCUSSION

The present study reports for the first time *in vivo* genotoxicity of combinations of Acm and Pro on sperm in germ cells of mice. The results of the present study revealed that Acm slightly, but not significantly, increased the percentage of abnormal sperm in mouse germ cells at all concentrations (0.625, 1.25 and 2.50) µg mL⁻¹ and exposure times (24 and 48 h). These findings suggest that Acm might not be a germ cell mutagen. According to the United States Environmental Protection Agency (US EPA) (29), Acm is classified as an “unlikely” human carcinogen. It has relatively low acute and chronic toxicity in mammals and there is no evidence of carcinogenicity, neurotoxicity, mutagenicity, and/or endocrine disruption. However, studies (8, 9) have reported genotoxic effects for Acm and other neonicotinoid insecticides (30-33). For instance, Bal et al. (34) reported that the neonicotinoid clothianidin induced impairment of the male mouse reproductive system. Such claims are inconsistent with the results of this study.

In the present study, Pro increased the percentage of abnormal sperm in mouse germ cells at the highest concentration (50 µg mL⁻¹) for 24 and 48 h. Quinto and De Marinis (35, 36) reported that Pro did not increase the percentage of abnormal sperm in mice. These reports are also in disagreement with our results.

However, the results of the present study support previous findings obtained by many other researchers. Prasad et al. (37) have reported that thiram increased the frequency of abnormal sperm in germ cells of mice. Hemavathi and Rahiman (38) stated that the commonly used dithiocarbamate fungicides ziram, thiram, and dithane M-45 induced a significant increase in the frequency of abnormal sperm at all tested doses. Giri et al. (39) have observed that the highest concentration of carbosulfan induced a >7-fold increase in the frequency of abnormal sperm. Ma et al. (40) found that the total sperm morphological abnormality was significantly higher in carbon disulphide-exposed workers.

There are no available investigations about the mutagenicity of Acm and Pro mixtures in germ cells of mice. Kocaman and Topaktas (9) have reported that the mixture of Acm and α-cypermethrin synergistically induced genotoxicity/cytotoxicity in human peripheral blood lymphocytes. Many researchers have found various pesticides, including Pro, to cause genotoxicity in peripheral blood lymphocytes of people applying pesticides in agriculture (2, 41-44). Furthermore, there are many investigations about the genotoxic effects of combinations of different pesticides (45-52).

For the sperm morphology assay, the treatment period had a wide time range due to the process of spermatogenesis. An understanding of the short-term, as well as the long-term, effects is essential if the causes of infertility and testicular atrophy are to be revealed cytogenetically. Many investigations evaluated effects of different chemicals using short-term exposure (14, 53-56). This was why, in the present study, 24 and 48 h applications were chosen for the short-term assay.

The exact reason of the increase in the frequency of abnormal sperm was not clear and opinions on this subject differ. The induction of abnormal sperms was assumed to be a result of an abnormal chromosome (57), minor alteration in testicular DNA (15), and point mutation (58). According to several studies (59-62), small deletions, point mutations, and abnormal chromosomes were proposed as possible genetic causes of such alterations. Bruce and Heddle (63) attributed the occurrence of sperm head abnormalities




Figure 5 Percentage of abnormal sperm (Mean±SE) in germ cells of mice exposed to mixtures of acetamiprid (*Acm*) and propineb (*Pro*) for 24 (a) and 48 h (b). Dark and white bars represent the measured values and the expected values, respectively. * and ** represent significant antagonistic and synergistic effects, respectively ($p < 0.05$)

to the chromosomal aberrations that occur during the packaging of genetic material in the sperm head or occurrence of point mutation in testicular DNA. Several studies (64-67) reported that abnormalities may also arise as a consequence of mistakes in the spermatozoa-differentiating process during spermatogenesis. Another two studies (66, 67) reported that abnormalities in sperm heads may occur by physiological, cytotoxic or genetic mechanisms or alterations in testicular DNA which in turn disrupts the process of differentiation of spermatozoa.

Conclusion and limitations

The results of the present study suggest that under the experimental conditions studied, *Acm* was non-genotoxic for mouse germ cells. *Pro*, however, may be a germ cell mutagen due to the observed increase in frequency of sperm abnormalities. Furthermore, higher concentrations of the *Acm* and *Pro* mixture antagonistically induced genotoxic effects of them on germ cells of mice. However, it also has to be mentioned that this study was limited by its use of commercially available pesticide formulations. Some of the effects observed in the present study might have been mediated by the other constituents of the pesticide formulations used. Although the biological activity of a pesticide is mostly determined by its active substance, the pesticide formulations available

on the market often consist of a pure active ingredient and different technical materials, which are usually used to improve the properties of a chemical for handling, storage, and application (68). In many cases, such materials contain various ingredients, some of which are protected by patents, which make it difficult to fully understand the results obtained with commercially available pesticides, as was the case in the present study.

Nevertheless, the findings of present study suggest that the sperm head abnormality assay may be used as a short-term and reliable biological indicator in the genotoxic bioassay of pesticides. However, to obtain better insight into the mutagenicity and DNA damaging potential of *Acm* and *Pro*, further studies at molecular level should be done.

Acknowledgements

This study was a part of a project entitled "Micronucleus induction in mice bone marrow and human lymphocytes after exposure to mixture of pesticides acetamiprid-propineb" and supported by the Duzce University Scientific Research Fund [Project Number = 2011.05.01.070], Turkey. The author thanks Meral Kekecoglu for support and Zekiye Kiris and Salih Tunc Kaya for laboratory assistance.

REFERENCES

1. Zhang WJ, Jiang FB, Ou JF. Global pesticide consumption and pollution: with China as a focus. *Proc Int Acad Ecol Environ Sci* 2011;1:125-44.
2. Pasquini R, Scassellati-Sforzolini G, Angeli G, Fatigoni C, Monarca S, Beneventi L, DiGiulio AM, Bauleo FA. Cytogenetic biomonitoring of pesticide-exposed farmers in central Italy. *J Environ Pathol Toxicol Oncol* 1996;15:29-39. PMID: 9037262
3. Santamaria AB, Gobbi M, Cembran M, Arnaboldi A. Human lymphocyte micronucleus genotoxicity test with mixtures of phytochemicals in environmental concentrations. *Mutat Res* 1997;388:27-32. PMID: 9025789
4. Lander F, Knudsen LE, Gamborg MO, Järventaus H, Norppa H. Chromosome aberrations in pesticide-exposed greenhouse workers. *Scand J Work Environ Health* 2000;26:436-42. doi: 10.5271/sjweh.565
5. Shaham J, Kaufman Z, Gurchik R, Levi Z. Frequency of sister-chromatid exchange among greenhouse farmers exposed to pesticides. *Mutat Res* 2001;491:71-80. PMID: 11287300
6. Costa C, Teixeira JP, Silva S, Roma-Torres J, Coelho P, Gaspar J, Alves M, Laffon B, Rueff J, Mayan O. Cytogenetic and molecular biomonitoring of a Portuguese population exposed to pesticides. *Mutagenesis* 2006;21:343-50. doi: 10.1093/mutage/gel039
7. Rasgele PG, Kaymak F. The cytogenetic effects of logran on bone marrow cells of *Mus musculus*. *Pakistan J Biol Sci* 2006;9:2781-6. doi: 10.3923/pjbs.2006.2781.2786
8. Kocaman AY, Topaktaş M. *In Vitro* evaluation of the genotoxicity of acetamiprid in human peripheral blood lymphocytes. *Environ Mol Mutagen* 2007;48:483-90. PMID: 17603792
9. Kocaman AY, Topaktaş M. Genotoxic effects of a particular mixture of acetamiprid and alpha-cypermethrin on chromosome aberration, sister chromatid exchange, and micronucleus formation in human peripheral blood lymphocytes. *Environ Toxicol* 2010;25:157-68. doi: 10.1002/tox.20485
10. Kaymak F, Rasgele PG. Genotoxic effects of raxil on root tips and anthers of *Allium cepa* L. *Caryologia* 2009;62:1-9.
11. Gökalp Muranlı FD, Güner U. Induction of micronuclei and nuclear abnormalities in erythrocytes of mosquito fish (*Gambusia affinis*) following exposure to the pyrethroid insecticide lambda-cyhalothrin. *Mutat Res* 2011;726:104-8. doi: 10.1016/j.mrgentox.2011.05.004
12. Kocaman AY, Rencüzoğulları E, Topaktaş M. *In vitro* investigation of the genotoxic and cytotoxic effects of thiacloprid in cultured human peripheral blood lymphocytes. *Environ Toxicol* 2012 [Epub ahead of print] doi: 10.1002/tox.21790
13. Sharpe RM. Declining sperm counts in men – is there an endocrine cause? *J Endocrinol* 1993;136:357-60. doi: 10.1677/joe.0.1360357
14. Hess RA. Effects of environmental toxicants on the efferent ducts, epididymis and fertility. *J Reprod Fertil* 1998;53(Suppl):247-59. PMID: 10645284
15. Giri S, Prasad SB, Giri A, Sharma GD. Genotoxic effects of malathion: an organophosphorus insecticide, using three mammalian bioassays *in vivo*. *Mutat Res* 2002;514:223-31. PMID: 11815260
16. Ieradi LA, Zima J, Allegra F, Kotlanova E, Campanella L, Grossi R, Cristaldi M. Evaluation of genotoxic damage in wild rodents from a polluted area in the Czech Republic. *Folia Zool* 2003;52:57-66.
17. Lu B, Wu X, Tie X, Zhang Y, Zhang Y. Toxicology and safety of anti-oxidant of bamboo leaves. Part 1: Acute and subchronic toxicity studies on anti-oxidant of bamboo leaves. *Food Chem Toxicol* 2005;43:783-92. PMID: 15778019
18. Wyrobek AJ, Gordon LA, Burkhart JG, Francis MW, Kapp RW Jr, Letz G, Malling HG, Topham JC, Whorton MD. An evaluation of the mouse sperm morphology test and other sperm tests in non-human mammals. A report of the United States Environmental Protection Agency Gene-Tox Programme. *Mutat Res* 1983;115:1-72. doi: 10.1016/0165-1110(83)90014-3
19. Casida JE, Quistad GB. Why insecticides are more toxic to insect than people: the unique toxicology of insects. *J Pestic Sci* 2004;29:81-6.
20. Tomizawa M, Yamamoto I. Structure-activity relationships of nicotinoids and imidacloprid analogs. *J Pestic Sci* 1993;18:91-8.
21. Barrera CH, Pardo LC, Cortina GD. Antracol WP 70 genotoxicity in human lymphocyte cultures. *Colomb Med* 2008;39(Suppl 2):29-34.
22. Soloneski S, Reigosa MA, Larramendy ML. Effect of the dithiocarbamate pesticide zineb and its commercial formulation azzurro. V. Abnormalities induced in the spindle apparatus of transformed and non-transformed mammalian cell lines. *Mutat Res* 2003;536:121-9. doi: 10.1016/S1383-5718(03)00038-X
23. Karaca C, Sahin ME, Turabi MS, Dursun N, Altunoglu CC, Bedir C, Akyazı H, Keles R, Bahce UU, Ozer O, Algan N, Ocalan G, Karatas P, Ocak Y. According to drug recommendations, Plant Protection Products, T. C. Ministry of Agriculture and Rural Affairs, General Directorate of Protection and Control, System Offset, Ankara, TR: 2009. P. 131, 238-39.
24. Rolandi A, De Marinis E, De Caterina M. Dithiocarbamate pesticides: Activity of Propineb in the micronucleus test in mice. *Mutat Res* 1984;135:193-7. PMID: 6424007
25. Çavaş T, Çinkılıç N, Vatan O, Yılmaz D, Coşkun M. *In vitro* genotoxicity evaluation of acetamiprid in CaCo-2 cells using the micronucleus, comet and gamma-H2AX foci assays. *Pest Biochem Physiol* 2012;104:212-7. doi: 10.1016/j.pestbp.2012.08.004
26. Bakare AA, Okunola AA, Adetunji OA, Jenmi HB. Genotoxicity assessment of a pharmaceutical effluent using four bioassays. *Genet Mol Biol* 2009;32:373-81. doi: 10.1590/S1415-47572009000200026
27. Wyrobek AJ, Bruce WR. Chemical induction of sperm abnormalities in mice. *Proc Natl Acad Sci USA* 1975;72:4425-9. doi: 10.1073/pnas.72.11.4425
28. Šegvić Klarić M, Rumora L, Ljubanović D, Pepeljnjak S. Cytotoxicity and apoptosis induced by fumonisin B1, beauvericin and ochratoxin A in porcine kidney PK15 cells: effects of individual and combined treatment. *Arch Toxicol* 2008;82:247-55. doi: 10.1007/s00204-007-0245-y
29. United States Environmental Protection Agency (US EPA). Office of Prevention, Pesticides and Toxic Substances (7501C), Pesticide Fact Sheet, Name of Chemical: Acetamiprid, Reason for Issuance: Conditional Registration 2002 [displayed 14 January 2014]. Available at <http://www.>

- epa.gov/pesticides/chem_search/reg_actions/registration/fs_PC-099050_15-Mar-02.pdf
30. Calderón-Segura ME, Gómez-Arroyo S, Villalobos-Pietrini R, Martínez-Valenzuela C, Carbajal-López Y, Calderón-Ezquerro Mdel C, Cortés-Eslava J, García-Martínez R, Flores-Ramírez D, Rodríguez-Romero MI, Méndez-Pérez P, Bañuelos-Ruiz E. Evaluation of genotoxic and cytotoxic effects in human peripheral blood lymphocytes exposed *in vitro* to neonicotinoid insecticides news. *J Toxicol* 2012;2012:612647. doi: 10.1155/2012/612647
 31. Zhang Y, Zhong Y, Luo Y, Kong ZM. Genotoxicity of two novel pesticides for the earthworm, *Eisenia fetida*. *Environ Pollution* 2000;108:271-8. PMID: 15092957
 32. Bal R, Naziroğlu M, Türk G, Yılmaz Ö, Kuloğlu T, Etem E, Baydas G. Insecticide imidacloprid induces morphological and DNA damage through oxidative toxicity on the reproductive organs of developing male rats. *Cell Biochem Funct* 2012;30:492-9. doi: 10.1002/cbf.2826
 33. Bal R, Türk G, Tuzcu M, Yılmaz O, Kuloğlu T, Gundogdu R, Gür S, Agca A, Ulas M, Cambay Z, Tuzcu Z, Gencoglu H, Guvenç M, Ozsahin AD, Kocaman N, Aslan A, Etem E. Assessment of imidacloprid toxicity on reproductive organ system of adult male rats. *J Environ Sci Health B* 2012;47:434-44. doi: 10.1080/03601234.2012.663311
 34. Bal R, Türk G, Yılmaz O, Etem E, Kuloğlu T, Baydas G, Naziroğlu M. Effects of clothianidin exposure on sperm quality, testicular apoptosis and fatty acid composition in developing male rats. *Cell Biol Toxicol* 2012;28:187-200. doi: 10.1007/s10565-012-9215-0
 35. Quinto I, De Marinis E. Effects of propineb and dowther A on sperm morphology in mice. *Mutat Res* 1982;97:215.
 36. Quinto I, De Marinis E. Evaluation of Propineb, a dithiocarbamate pesticide, in the mouse-sperm morphology assay. *Mutat Res* 1983;124:235-40. PMID: 6656825
 37. Prasad MH, Pushpavathi K, Rita P, Reddy PP. The effect of thiram on the germ cells of male mice. *Food Chem Toxicol* 1987;25:709-11. PMID: 3653825
 38. Hemavathi E, Rahiman MA. Toxicological effects of ziram, thiram, and dithane M-45 assessed by sperm shape abnormalities in mice. *J Toxicol Environ Health* 1993;38:393-8. PMID: 8478981
 39. Giri S, Giri A, Sharma GD, Prasad SB. Mutagenic effects of carbosulfan, a carbamate pesticide. *Mutat Res* 2002;519:75-82. PMID: 12160893
 40. Ma JY, Ji JJ, Ding Q, Liu WD, Wang SQ, Wang N, Chen GY. The effects of carbon disulfide on male sexual function and semen quality. *Toxicol Ind Health* 2010;26:375-82. doi: 10.1177/0748233710369127
 41. Bolognesi C, Parrini M, Merlo F, Bonassi S. Frequency of micronuclei in lymphocytes from a group of floriculturists exposed to pesticides. *J Toxicol Environ Health* 1993;40:405-11. PMID: 8230311
 42. Falck GC, Hirvonen A, Scarpato R, Saarikoski ST, Migliore L, Norppa H. Micronuclei in blood lymphocytes and genetic polymorphism for GSTM1, GSTT1 and NAT2 in pesticide-exposed greenhouse workers. *Mutat Res* 1999;441:225-37. PMID: 10333536
 43. Pastor S, Gutiérrez S, Creus A, Xamena N, Piperakis S, Marcos R. Cytogenetic analysis of Greek farmers using the micronucleus assay in peripheral lymphocytes and buccal cells. *Mutagenesis* 2001;16:539-45. doi: 10.1093/mutage/16.6.539
 44. Paz-y-Miño C, Bustamante G, Sánchez ME, Leone PE. Cytogenetic monitoring in a population occupationally exposed to pesticides in Ecuador. *Environ Health Perspect* 2002;110:1077-80. PMID: 12417477
 45. Dolara P, Salvadori M, Capobianco T, Torricelli F. Sister chromatid exchanges in human lymphocytes induced by dimethoate, omethoate, deltamethrin, benomyl and their mixture. *Mutat Res* 1992;283:113-8. PMID: 1381487
 46. Roloff DB, Belluck AD, Meisner FL. Cytogenetic studies of herbicide interactions *in vitro* and *in vivo* using atrazine and linuron. *Arch Environ Contam Toxicol* 1992;22:267-71. doi: 10.1007/BF00212084
 47. Rencüzoğullari E, Topaktaş M. Sister chromatid exchange in cultured human lymphocytes treated with carbosulfan, ethyl carbamate and ethyl methanesulfonat separately and in mixtures. *Tr J Biol* 1998;22:369-87.
 48. Rencüzoğullari E, Topaktaş M. Chromosomal aberrations in cultured human lymphocytes treated with the mixtures of carbosulfan, ethyl carbamate and ethyl methanesulfonate. *Cytologia* 2000;65:83-92. doi: 10.1508/cytologia.65.83
 49. Chauhan LKS, Chandra S, Saxena PN, Gupta SK. *In vivo* cytogenetic effects of a commercially formulated mixture of cypermethrin and quinalphos in mice. *Mutat Res* 2005;587:120-5. PMID:16185912
 50. Karabay UN, Oguz MG. Cytogenetic and genotoxic effects of the insecticides, imidacloprid and methamidophos. *Gen Mol Res* 2005;4:653-62. PMID: 16475109
 51. Demsia G, Vlastos D, Goumenou M, Demetrios PA. Assesment of the genotoxicity of imidacloprid and metalaxyl in cultured human lymphocytes and rat bone-marrow. *Mutat Res* 2007;634:32-9. doi: 10.1016/j.mrgentox.2007.05.018
 52. Das PP, Shaik AP, Jamil K. Genotoxicity induced by pesticide mixtures: *in-vitro* studies on human peripheral blood lymphocytes. *Toxicol Ind Health* 2007;7:449-58. doi: 10.1177/0748233708089040
 53. Dobrzyńska MM, Gajewski AK. Induction of micronuclei in bone marrow and sperm head abnormalities after combined exposure of mice to low doses of X-rays and acrylamide. *Teratog Carcinog Mutagen* 2000;20:133-40. PMID: 10820423
 54. Latchoumycandane C, Chitra KC, Mathur PP. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) induces oxidative stress in the epididymis and epididymal sperm of adult rats. *Arch Toxicol* 2003;77:280-4. PMID: 12734642
 55. Chandra P, Khuda-Bukhsh AR. Genotoxic effects of cadmium chloride and azadirachtin treated singly and in combination in fish. *Ecotoxicol Environ Saf* 2004;58:194-201. PMID: 15157573
 56. Uzunhisarcikli M, Kalender Y, Dirican K, Kalender S, Ogutcu A, Buyukkomurcu F. Acute, subacute and subchronic administration of methyl parathion-induced testicular damage in male rats and protective role of vitamins C and E. *Pestic Biochem Physiol* 2007;87:115-22.
 57. Bruce WR, Furrer R, Wyrobek AJ. Abnormalities in the shape of murine sperm after acute testicular X-irradiation. *Mutat Res* 1974;23:381-6. PMID: 4407800
 58. Narayana K, D'Souza UJ, Seetharama Rao KP. Ribavirin-induced sperm shape abnormalities in Wistar rat. *Mutat Res* 2002;513:193-6. PMID: 11719104
 59. Mendoza-Lujambio I, Burfeind P, Dixkens C, Meinhardt A, Hoyer-Fender S, Engel W, Neesen J. The Hook1 gene is non-functional in the abnormal spermatozoon head shape

- (azh) mutant mouse. *Human Mol Genet* 2002;11:1647-58. doi: 10.1093/hmg/11.14.1647
60. Pyle A, Handel MA. Meiosis in male PL/J mice: a genetic model for gametic aneuploidy. *Mol Reprod Dev* 2003;64:471-81. doi: 10.1002/mrd.10231
61. Escalier D, Bai XY, Silviu D, Xu PX, Xu X. Spermatid nuclear and sperm periaxonemal anomalies in the Mouse *Ube2b* null mutant. *Mol Reprod Dev* 2003;65:298-308. doi: 10.1002/mrd.10290
62. Lamara AS, Fonseca G, Fuentes JL, Cozzi R, Cundari E, Fiore M, Ricordy R, Perticone P, Degrassi F, De Salvia R. Assessment of the genotoxic risk of *Punica granatum* L. (Punicaceae) whole fruit extracts. *J Ethnopharmacol* 2008;115:416-22. PMID: 18036756
63. Bruce W, Heddle J. The mutagenicity of 61 agents as determined by the micronucleus, Salmonella and sperm abnormality assays. *Can J Cytol Genet* 1979;21:319-34. PMID: 393369
64. Bakare AA, Mosuro AA, Osibanjo O. An *in vivo* evaluation of induction of abnormal sperm morphology in mice by land fill leachates. *Mutat Res* 2005;582:28-34. PMID: 15781207
65. Otitoloju AA, Obe IA, Adewale OA, Otubanjo OA, Osunkalu VO. Preliminary study on the induction of sperm head abnormalities in mice, *Mus musculus*, exposed to radiofrequency radiations from global system for mobile communication base stations. *Bull Environ Contam Toxicol* 2010;84:51-4. doi: 10.1007/s00128-009-9894-2
66. Odeigah PGC, Nurudeen O, Amund OO. Genotoxicity of oil field wastewater in Nigeria. *Hereditas* 1997;126:161-8. doi: 10.1111/j.1601-5223.1997.00161.x
67. Agunbiade SO, Okonko IO, Alimba CG, Folarin AC, Anugweje KC. Effects of a carbonaceous bottling plant effluent on albino mice sperm morphology and testes histopathology. *Nat Sci* 2012;10:154-60 [displayed 10 May 2013]. Available at http://www.sciencepub.net/nature/ns1008/024_10572ns1008_154_160.pdf
68. Knowles DA. *Chemistry and Technology of Agrochemical Formulations*. London: Kluwer Academic Publishers; 1998

Sažetak

Abnormalna morfologija mišjih spermija nakon kratkotrajne izloženosti acetamipridu, propinebu i njihovoj mješavini

Pesticidi su snažni zagađivači okoliša, a akumuliraju se i u biotičkim i abiotičkim sastavnicama ekosustava. Acetamiprid (Acm), insekticid iz skupine neonikotinoidea, i propineb (Pro), fungicid iz skupine ditiokarbamata, imaju široku primjenu u kontroli sisajućih insekata, odnosno gljivičnih infekcija, na usjevima. Ovim se radom istražuju genotoksični učinci spomenutih spojeva, pojedinačno i u mješavini, na mišje spolne stanice testom morfološke analize spermija. Miševima je intraperitonealno ubrizgano 0,625, 1,25 i 2,50 $\mu\text{g mL}^{-1}$ Acm-a, 12,5, 25,00 i 50,00 $\mu\text{g mL}^{-1}$ Pro-a, te njihova mješavina pri istim koncentracijama tijekom 24 i 48 sati. Acm nije povećao postotak abnormalnih spermija ni pri jednoj koncentraciji. Učestalost abnormalnih spermija značajno je porasla nakon 24 i 48 sati izloženosti 50 $\mu\text{g mL}^{-1}$ Pro-a. Mješavina od 2,50 $\mu\text{g mL}^{-1}$ Acm-a i 50,00 $\mu\text{g mL}^{-1}$ Pro-a uzrokovala je abnormalnosti spermija antagonistički i nakon 24 i nakon 48 sati izloženosti. Naši rezultati pokazuju kako Acm nije bio genotoksičan za mišje spolne stanice, a primijećeni porast učestalosti abnormalnosti spermija nakon izlaganja Pro upućuje na to da bi on mogao biti mutagen. Međutim, kako bi se stekao bolji uvid u mutagenost i potencijal obaju pesticida za oštećenje DNA, nužna su daljnja istraživanja na molekularnoj razini.

KLJUČNE RIJEČI: kombinirani učinci; mješavina pesticida; test morfološke analize spermija

CORRESPONDING AUTHOR:

Pinar G. Rasgele
Duzce University, Faculty of Agriculture and Natural
Sciences, Department of Biosystems Engineering
81620, Duzce, Turkey
E-mail: pinargocrasgele@gmail.com