

## INDOOR PM<sub>1</sub>, PM<sub>2.5</sub>, PM<sub>10</sub> AND OUTDOOR PM<sub>2.5</sub> CONCENTRATIONS IN PRIMARY SCHOOLS IN SARI, IRAN

Mahmoud MOHAMMADYAN<sup>1</sup> and Bijan SHABANKHANI<sup>2</sup>

*Health Sciences Research Center<sup>1</sup>, Faculty of Health<sup>2</sup>, Mazandaran University of Medical Sciences, Sari, Iran*

Received in January 2013  
CrossChecked in April 2013  
Accepted in July 2013

This study was carried out to determine the distribution of particles in classrooms in primary schools located in the centre of the city of Sari, Iran and identify the relationship between indoor classroom particle levels and outdoor PM<sub>2.5</sub> concentrations. Outdoor PM<sub>2.5</sub> and indoor PM<sub>1</sub>, PM<sub>2.5</sub>, and PM<sub>10</sub> were monitored using a real-time Micro Dust Pro monitor and a GRIMM monitor, respectively. Both monitors were calibrated by gravimetric method using filters. The Kolmogorov-Smirnov test showed that all indoor and outdoor data fitted normal distribution. Mean indoor PM<sub>1</sub>, PM<sub>2.5</sub>, PM<sub>10</sub> and outdoor PM<sub>2.5</sub> concentrations for all of the classrooms were 17.6 µg m<sup>-3</sup>, 46.6 µg m<sup>-3</sup>, 400.9 µg m<sup>-3</sup>, and 36.9 µg m<sup>-3</sup>, respectively. The highest levels of indoor and outdoor PM<sub>2.5</sub> concentrations were measured at the Shahed Boys School (69.1 µg m<sup>-3</sup> and 115.8 µg m<sup>-3</sup>, respectively). The Kazemi school had the lowest levels of indoor and outdoor PM<sub>2.5</sub> (29.1 µg m<sup>-3</sup> and 15.5 µg m<sup>-3</sup>, respectively). In schools located near both main and small roads, the association between indoor fine particle (PM<sub>2.5</sub> and PM<sub>1</sub>) and outdoor PM<sub>2.5</sub> levels was stronger than that between indoor PM<sub>10</sub> and outdoor PM<sub>2.5</sub> levels. Mean indoor PM<sub>2.5</sub> and PM<sub>10</sub> and outdoor PM<sub>2.5</sub> were higher than the standards for PM<sub>2.5</sub> and PM<sub>10</sub>, and there was a good correlation between indoor and outdoor fine particle concentrations.

**KEY WORDS:** *indoor particle concentrations, outdoor particle concentrations*

Recent epidemiological studies have documented an association between changes in ambient particulate matter (PM) concentrations and changes in daily mortality and morbidity (1-3). Furthermore, air quality at schools seems to be a major determinant of health outcomes (4). Most of these studies have emphasised the importance of particulate matter with aerodynamic diameter of less than 10 µm (PM<sub>10</sub>) and, recently, 2.5 µm (PM<sub>2.5</sub>), measured at fixed monitoring sites. It seems that traffic-related particles are more toxic than others. Peters et al. (5) have reported that the risk of exposure to black carbon (BC) as a surrogate of traffic particles is higher than to other ambient particulates (5). Schwarz et al. (6) have also reported a stronger association of BC than of PM<sub>2.5</sub> with changes in heart

rate. In a multi-city study, Dominci et al. (7) have shown that the association between PM<sub>10</sub> concentrations and increased risk of death generally remains unchanged after control for other air pollutants.

People spend considerable time indoors: at home, school, work or in vehicles (8-10). School children, the elderly, and other groups of people more susceptible to the effects of poor air quality spend even more time indoors (11). Several studies have reported high concentrations of PM in classrooms (12-17). Major studies about personal exposure to particles (18, 19) have found poor correlation between personal exposure to fine particulate matter and outdoor air particle concentrations, but they have also reported

good correlation between personal exposure and indoor air particle concentrations.

The aim of this study was to address this issue by determining the distribution of  $PM_{10}$ ,  $PM_{2.5}$ , and  $PM_{10}$  in the classrooms of primary schools located in the centre of the city of Sari and by identifying the relationship between indoor and outdoor  $PM_{2.5}$  concentrations.

Sari is the capital of the Iranian province of Mazandaran, located some 30 km to the south of the Caspian Sea and stretching from the northern slopes of the Elburz Mountains to across the Tajan River. It has a population of 270,000 people residing in the town and about as many residing in the suburbs. Our earlier reports have shown that personal exposure to  $PM_{10}$  among taxi and bus drivers and to  $PM_{2.5}$  in shops in the city centre area are higher than the standards recommended by the US Environmental Protection Agency (EPA) (20-22).

## MATERIALS AND METHODS

Our indoor and outdoor monitoring involved primary schools located in the centre of Sari with four major roads of varying traffic density. In Enghlab Street (south) it is about 1,500 vehicles per hour, in 18-Day Street (east) 2,220, in Jomhori Street (west) 1,260, and in Modarres Street (south) 1,250. Khosravi School is located in Enghlab Street; Shahed Boy and Shahed Girl Schools are located on Modarres Street; Kazemi School, Ghaemi School, and Ameneh School are located on three smaller roads less than 100 m away from Jomhori Street. School buildings are about 10 to 40 years old. Classrooms - all accommodating between 27 and 32 pupils - have a similar design, and their area varies from 24 m<sup>2</sup> to 34.2 m<sup>2</sup>. Floors are stone. No mechanical ventilation or air conditionings were in use during the monitoring period. However, all classrooms were heated by radiators in the cold months.

Over 26 days of a school year (spanning from November 2011 to June 2012), we monitored  $PM_{10}$ ,  $PM_{2.5}$ , and  $PM_{10}$  concentrations indoors and  $PM_{2.5}$  concentrations outdoors. Both indoor and outdoor monitoring started and ended with the classes (from 8:00 a.m. to around 12:30 p.m.). Average indoor monitoring time was 4.39 h (range 2.95 h to 4.7 h), depending on the duration of a particular class. The indoor dust monitor was placed in the centre of the classroom, about 80 cm above the floor, and the

outdoor monitor in the school yard at least one metre away from any obstacle and one metre above ground.

For indoor measurements we used a GRIMM real-time aerosol spectrometer and dust monitor (Model 1.108, Grimm Aerosol Technik GmbH, Ainring, Germany).  $PM_{10}$ ,  $PM_{2.5}$  and  $PM_{10}$  concentrations were recorded at one-minute intervals. This dust aerosol spectrometer has been designed for continuous particle count and for calculating particle mass based on particle density. It has an integrated gravimetric filter that collects all particles after optical measurement for further analysis. Data can be displayed as particle concentration and as mass concentration. Sample air is sucked through a measuring cell and a gravimetric filter by an internal flow-control pump. The filter serves as a dust collector and as gravimetric control of optical measurements.

For outdoor measurements we used a MicroDust Pro real-time monitor (Casella, Bedford, UK). This instrument is calibrated to a known reference dust standard. Different dust types cause a different response from this instrument due to variation in particle size, refractive indices, particle density, and colour. In order to correct for this, it is necessary to calibrate the response of the instrument. This involves the collection of a gravimetric (filtered) sample of the dust after it has passed through the probe optics. To measure  $PM_{2.5}$  concentrations, a size-selective sampling cyclone was used in combination with a particle size adaptor and a small polyurethane foam (PUF) filter that was designed for  $PM_{2.5}$  size fraction monitoring. A small personal sampling pump was used to provide continuous air flow through the gravimetric adaptor and photo detector. For gravimetric calibration, particles were then collected on a 37 mm, 2.0  $\mu$ m Teflon filter (SKC Inc., Dorset, UK), which was placed in the cassette behind air sample stream. To obtain mean  $PM_{2.5}$  concentrations we divided particle mass (in  $\mu$ g), obtained by weighing the filter, with the volume of sampled air drawn through the instrument (in m<sup>3</sup>). For calibration we compared mean  $PM_{2.5}$  concentration with the average  $PM_{2.5}$  concentration obtained from direct reading from the MicroDust Pro instrument.

The results for each location had to be corrected with a gravimetric factor – the so-called C-factor. To determine the C-factor and to compare the displayed data, the GRIMM dust monitor and the MicroDust Pro monitor were run side by side in six classrooms for five hours, one day a month over the study period.

The GRIMM monitor was run on the particle concentration mode to measure particles between 0.3  $\mu\text{m}$  and 20  $\mu\text{m}$ , and the MicroDust monitor was run to measure  $\text{PM}_{2.5}$ . Filters were desiccated for 24 h and weighed with a microbalance (resolution 1  $\mu\text{g}$ ) three times before and after sampling. Total dust weight on filters was divided with the calculated total volume of air sucked by pumps to determine mean gravimetric concentrations of particles. Running both instruments side by side provided information on actual average gravimetric concentrations, which were then divided by mean particle concentrations downloaded from respective instruments to obtain gravimetric calibration factors. Finally, all real-time data were multiplied by calibration factors obtained for either instrument to obtain actual particle concentrations. In total, we collected data for 7,115 one-minute indoor and outdoor particle concentration readings. Mean correction factors of 1.03 and 1.14 were applied for the GRIMM and the MicroDust Pro monitor data, respectively. One-minute data were used for statistical analysis.

#### Statistical analysis

The statistic package SPSS v.17 for windows was used for running the Kolmogorov-Smirnov test (K-S test) to assess the normality of the frequency distributions of  $\text{PM}_1$ ,  $\text{PM}_{2.5}$ , and  $\text{PM}_{10}$  concentrations. This statistic package also was used for running descriptive statistics and univariate regression model to assess the association between outdoor  $\text{PM}_{2.5}$  concentrations and indoor classroom  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$ , and  $\text{PM}_1$ . The Microsoft Office EXCEL 2007 software was

used to make a graph for demonstration of daily mean indoor  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  concentrations.

## RESULTS AND DISCUSSION

The Kolmogorov-Smirnov (K-S) test shows that all indoor particle concentration data fit normal distribution (Figures 1-3). The indoor classroom  $\text{PM}_{2.5}$  concentrations show distributions that are bi-modal, suggesting that there may be outliers within the indoor classroom  $\text{PM}_{2.5}$  data. Resuspension of fine particles as a result of student activities may explain transient high indoor classroom particle concentrations. Mean indoor  $\text{PM}_1$ ,  $\text{PM}_{2.5}$ ,  $\text{PM}_{10}$ , and outdoor  $\text{PM}_{2.5}$  concentrations were 17.6  $\mu\text{g m}^{-3}$ , 46.6  $\mu\text{g m}^{-3}$ , 400.9  $\mu\text{g m}^{-3}$ , and 36.9  $\mu\text{g m}^{-3}$  respectively. Figure 4 shows daily mean indoor classroom  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  levels. On some days, mean  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  exceeded the respective US EPA standards of 35  $\mu\text{g m}^{-3}$  and 150  $\mu\text{g m}^{-3}$  (23). High concentrations of  $\text{PM}_{10}$  could be due to resuspension of chalk dust, skin flaks, and insect dander that can increase the particle concentration when the students are active. In Tehran, Halek et al. (17) reported mean indoor classroom  $\text{PM}_1$ ,  $\text{PM}_{2.5}$ , and  $\text{PM}_{10}$  of 19  $\mu\text{g m}^{-3}$ , 42  $\mu\text{g m}^{-3}$ , and 274  $\mu\text{g m}^{-3}$ , respectively. Mean  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  concentrations were lower than in our study (46.6  $\mu\text{g m}^{-3}$  and 400.9  $\mu\text{g m}^{-3}$ , respectively), whereas mean indoor  $\text{PM}_1$  concentration and mean outdoor  $\text{PM}_{2.5}$  in our study were similar with those reported by in Tehran (17.6  $\mu\text{g m}^{-3}$  and 36.9  $\mu\text{g m}^{-3}$  vs. 19  $\mu\text{g m}^{-3}$  and 38  $\mu\text{g m}^{-3}$ , respectively). In Munich, Germany,




Figure 1 Distribution of indoor classroom  $\text{PM}_1$  concentrations






Figure 2 Distribution of indoor classroom  $\text{PM}_{2.5}$  concentrations



**Figure 3** Distribution of indoor classroom  $PM_{10}$  concentrations

Fromme et al. (15) reported lower indoor and outdoor median  $PM_{2.5}$  and  $PM_{10}$  than we have, but our findings are lower than those reported for Athens and Istanbul (25, 26).



**Figure 4** Mean indoor classroom  $PM_{2.5}$  and  $PM_{10}$  concentrations

Table 1 shows the descriptive statistics for indoor  $PM_{10}$ ,  $PM_{2.5}$ ,  $PM_{10}$  and outdoor  $PM_{2.5}$  in monitored schools. Khosravi school showed the highest mean  $PM_{10}$  concentration and Ameneh School the lowest. The highest indoor and outdoor  $PM_{2.5}$  concentration was recorded in Shahed Boy School ( $69.1 \mu\text{g m}^{-3}$  and  $115.8 \mu\text{g m}^{-3}$ , respectively). Kazemi School, in turn, showed the lowest indoor and outdoor  $PM_{2.5}$  ( $29.1 \mu\text{g m}^{-3}$  and  $15.5 \mu\text{g m}^{-3}$ , respectively). Despite

**Table 1** Outdoor  $PM_{2.5}$  and indoor classroom  $PM_{10}$ ,  $PM_{2.5}$  and  $PM_1$  concentrations by school

Location	School	N	Mean	SD	Min	Max
Indoor $PM_{10}$	Khosravi	1284	568.2	353.2	44.7	2911.1
	Kazemi	1233	286.9	129.9	55.2	898.6
	Ghaemi	1064	381.0	275.8	23.9	2036.4
	Ameneh	1355	265.8	157.6	62.9	994.5
	Shahed boys	1078	428.2	261.3	68.9	1752.5
	Shahed girls	1101	492.5	503.6	54.6	5858.5
Indoor $PM_{2.5}$	Khosravi	1284	49.8	22.6	10.4	136.9
	Kazemi	1233	29.1	16.8	6.5	156.8
	Ghaemi	1064	38.2	20.9	6.2	182.7
	Ameneh	1355	38.3	16.3	6.2	87.9
	Shahed boys	1078	69.1	36.5	9	141.1
	Shahed girls	1101	59.0	62.0	6.7	504.7
Indoor $PM_1$	Khosravi	1284	12.1	4.2	3.7	28.9
	Kazemi	1233	14.1	14.5	2.8	140.1
	Ghaemi	1064	14.9	12.0	3.9	151.3
	Ameneh	1355	21.0	10.5	2.4	50
	Shahed boys	1078	28.9	15.2	3.1	51.7
	Shahed girls	1101	15.6	14.6	2.6	99.6
Outdoor $PM_{2.5}$	Khosravi	1284	22.1	23.1	0	196.1
	Kazemi	1233	15.5	18.4	0	201.8
	Ghaemi	1064	26.2	48.1	0	963.7
	Ameneh	1355	25.9	28.3	0	138.8
	Shahed boys	1078	115.8	146.9	1.2	498.2
	Shahed girls	1101	24.938	25.3	0	88.1

**Table 2** Relationship between outdoor  $PM_{2.5}$  and indoor classroom  $PM_{10}$ ,  $PM_{2.5}$ , and  $PM_1$  levels in schools located close to main or small roads.

PM	School location	Indoor $PM_{10}$		Indoor $PM_{2.5}$		Indoor $PM_1$	
		R	Sig	R	Sig.	R	Sig.
Outdoor $PM_{2.5}$	Main roads	0.09	0.00	0.35	0.00	0.51	0.00
	Small roads	0.08	0.00	0.30	0.00	0.62	0.00

the highest average  $PM_{10}$ , Khosravi School also had the lowest mean  $PM_1$  concentration ( $12.1 \mu\text{g m}^{-3}$ ).

We established a significant correlation between indoor classroom  $PM_1$ ,  $PM_{2.5}$ , and  $PM_{10}$  concentrations and outdoor  $PM_{2.5}$  concentrations for both schools located near main and small roads. However, Shahed boys school, which located on a main road, showed the highest mean indoor and outdoor  $PM_{2.5}$  levels and Kazemi school located on a small road had a lowest mean indoor and outdoor  $PM_{2.5}$ . Mean indoor  $PM_{10}$  and  $PM_{2.5}$  concentrations in schools located on the main roads were significantly higher than in schools on small roads ( $486.3 \mu\text{g m}^{-3}$  and  $49.2 \mu\text{g m}^{-3}$  vs.  $320.7 \mu\text{g m}^{-3}$  and  $44.3 \mu\text{g m}^{-3}$ , respectively). However, mean indoor  $PM_1$  concentration was higher in schools on small roads than in schools on the main roads ( $21.0 \mu\text{g m}^{-3}$  vs.  $14.0 \mu\text{g m}^{-3}$  respectively). One possible explanation is that  $PM_1$  and  $PM_{2.5}$ , which were mainly emitted from combustion sources, can distribute in the ambient easily and might be an effective factor for indoor particle concentrations. Similar studies concluded that proximity to traffic is a major determinant of the level of student exposure (12, 27).

The association between indoor fine particle concentrations ( $PM_{2.5}$  and  $PM_1$ ) and outdoor  $PM_{2.5}$  levels was stronger than between indoor  $PM_{10}$  concentrations and outdoor  $PM_{2.5}$  levels (Table 2). This is most likely because outdoor air enters classrooms through doors and windows, which are generally left open because of a moderate climate in Sari. On the other hand, the reason for the weak correlation between coarse  $PM_{10}$  indoor particles and outdoor  $PM_{2.5}$  concentrations is that coarse particles are mainly produced by indoor student activities. In contrast to our study, in which indoor  $PM_{2.5}$  was higher than outdoor  $PM_{2.5}$ , Buonanno et al. (24) reported higher concentrations of outdoor than indoor particles of up to  $3 \mu\text{m}$  in Cassino, Italy (24).

#### Acknowledgment

We wish to thank all primary school managers and teachers for their cooperation in the study. Our thanks

also goes to Mazandaran University of Medical Sciences for financial support.

#### REFERENCES

- Dockery DW, Pope CA III. Acute respiratory effects of particulate air pollution. *Annu Rev Public Health* 1994;15:107-32. doi: 10.1146/annurev.pu.15.050194.000543
- Seaton A, MacNee W, Donaldson K, Godden D. Particulate air pollution and acute health effects. *Lancet* 1995;345:176-8. doi: 10.1016/S0140-6736(95)90173-6
- Wallace LA. Correlations of personal exposure to particles with outdoor air measurements: a review of recent studies. *Aerosol Sci Technol* 2000;32:15-25. doi: 10.1080/027868200303894
- Mejia JF, Choy SL, Mengersen K, Morawska L. Methodology for assessing exposure and impacts of air pollutants in school children: Data collection, analysis and health effects – A literature review. *Atmos Environ* 2011;45:813-23. doi: 10.1016/j.atmosenv.2010.11.009
- Peters A, Dockery DW, Muller JE, Mittleman MA. Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 2001;103:2810-5. doi: 10.1161/01.CIR.103.23.2810
- Schwartz J, Litonjua A, Suh H, Verrier M, Zanobetti A, Syring M, Nearing B, Verrier R, Stone P, MacCallum G, Speizer FE, Gold DR. Traffic related pollution and heart rate variability in a panel of elderly subjects. *Thorax* 2005;60:455-61. doi: 10.1136/thx.2004.024836
- Dominici F, McDermott A, Daniels M, Zeger SL, Samet JM. Revised analyses of the National Morbidity, Mortality, and Air Pollution Study: mortality among residents of 90 cities. *J Toxicol Environ Health A* 2005;68:1071-92. doi: 10.1080/15287390590935932
- Mohammadyan M, Ashmore MR. Personal exposure and indoor  $PM_{2.5}$  concentrations in an urban population. *Indoor Built Environ* 2005;14:313-20. doi: 10.1177/1420326X05054293
- Wallace LA. Indoor particles: a review. *J Air Waste Manag Assoc* 1996;46:98-126. PMID: 8846246
- Spengler JD, Sexton K. Indoor air pollution: a public health perspective. *Science* 1983;221:9-17. doi: 10.1126/science.6857273
- Klepeis NE, Nelson WC, Ott WR, Robinson JP, Tsang AM, Switzer P, Behar JV, Hern SC, Engelmann WH. The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants. *J Exposure Anal Environ Epidemiol* 2001;11:231-52. PMID: 11477521

12. Janssen NAH, van Vliet PHN, Aaarts F, Harssema H, Brunekreef B. Assessment of exposure to traffic related air pollution of children attending schools near motorways. *Atmos Environ* 2001;35:3875-84. doi: 10.1016/S1352-2310(01)00144-3
13. Link B, Gabrio T, Zöllner I, Schwenk M, Siegel D, Schultz E, Scharring S, Borm P. [Programm Lebensgrundlage Umwelt und ihre Sicherung (BWPLUS) Forschungsbericht FZKA-BWPLUS, Feinstaubbelastungen und deren gesundheitliche Wirkungen bei Kindern, in German] 2004 [displayed 10 Juni 2013]. Available at <http://www.fachdokumente.lubw.baden-wuerttemberg.de/servlet/is/40191/BWB21007Sber.pdf?command=downloadContent&filename=BWB21007Sber.pdf>
14. Son BS, Song MR, Yang WH. A Study on PM<sub>10</sub> and VOCs concentrations of indoor environment in school and recognition of indoor air quality. In: from 10th International Conference on Indoor Air Quality and Climate; 4-9 Sep 2005. Beijing, China. Beijing: Tshinghua University Press; 2005. p. 827-32.
15. Fromme H, Dietrich S, Twardella D, Heitmann D, Schierl R, Liebl B, Rüden H. Particulate matter in the indoor air of classrooms - exploratory results from Munich and surrounding. *Atmos Environ* 2007;41:854-66. doi: 10.1016/j.atmosenv.2006.08.053
16. Stranger M, Potgieter-Vermaak SS, Van Grieken R. Comparative overview of indoor air quality in Antwerp, Belgium. *Environ Int* 2007;33:789-97. doi: 10.1016/j.envint.2007.02.014
17. Halek F, Kianpour-rad M, Kavousirahim A. Parametric evaluation of indoor particulate matter in elementary schools in the central parts of Tehran. *Indoor Built Environ* 2013; 22:580-585. doi: 10.1177/1420326X11433224
18. Koistinen JK, Otto H, Tuulia R, Edwards RD, Moschandreas D, Jantunen MJ. Behavioural and Environmental determinants of personal exposure to PM<sub>2.5</sub> in EXPOLIS – Helsinki Finland. *Atmos Environ* 2001; 35:2473-2481. Doi:10.1177/1420326X11421510
19. Adgate JL, Ramachandran G, Pratt GC, Waller LA, Sexton K. Longitudinal variability in outdoor, indoor and personal PM<sub>2.5</sub> exposure in healthy non smoking adults. *Atmos Environ* 2003; 37:993-1002. Doi:10.1177/1420326X11421510
20. Mohammadyan M, Alizadeh A, Mohammadpour RA. Personal exposure to PM<sub>10</sub> among bus drivers in Sari, Iran. *Indoor Built Environ* 2009;18:83-9. doi: 10.1177/1420326X08101530
21. Mohammadyan M, Alizadeh A, Etemadinejad S. Personal exposure to PM<sub>10</sub> among taxi drivers in Iran. *Indoor Built Environ* 2010;19:538-45. doi: 10.1177/1420326X10378802
22. Mohammadyan M, Sojodi L, Etemadinejad S. Survey of concentrations of PM<sub>2.5</sub> indoor and outdoor of shops in Sari city center. *J Mazand Univ Med Sci* 2011;21:72-9.
23. U.S. Environmental Protection Agency (US EPA). National Ambient Air Quality Standards (NAAQS) [displayed 20 Sep 2012]. Available at <http://www.epa.gov/air/criteria.html>
24. Buonanno G, Fuoco FC, Morawska L, Stabile L. Airborne particle concentrations at school measured at different spatial scale. *Atmos Environ* 2013;67:38-45. doi: 10.1016/j.atmosenv.2012.10.048
25. Diapouli E, Chaloulakou A, Mihalopoulos N, Spyrellis N. Indoor and outdoor PM mass and number concentrations at schools in the Athens area. *Environ Monit Assess* 2008;136:13-20. doi: 10.1007/s10661-007-9724-0
26. Ekmekcioglu D, Keskin SS. Characterization of indoor air particulate matter in selected elementary schools in Istanbul, Turkey. *Indoor Built Environ* 2007;16:169-76. doi: 10.1177/1420326X07076777
27. Van Roosbroeck S, Jacobs J, Janssen NAH, Oldenwening M, Hoek G, Brunekreef B. Long-term personal exposure to PM<sub>2.5</sub>, soot and NO<sub>x</sub> in children attending schools located near busy roads, a validation study. *Atmos Environ* 2007;41:3381-94. doi: 10.1016/j.atmosenv.2006.12.023

## Sažetak

### KONCENTRACIJE LEBDEĆIH ČESTICA $PM_1$ , $PM_{2,5}$ , $PM_{10}$ U ZATVORENOM PROSTORU TE KONCENTRACIJE $PM_{2,5}$ ČESTICA U OTVORENOM PROSTORU OSNOVNIH ŠKOLA U GRADU SARIJU U IRANU

Svrha je ovog istraživanja bila utvrditi raspodjelu lebdećih čestica u osnovnim školama u središtu iranskoga grada Sarija te vidjeti jesu li razine lebdećih čestica mjenjenih u dvorištima škola i u učionicama međusobno povezane. Vani su mjerene  $PM_{2,5}$  čestice pomoću stalnog Micro Dust Pro monitora, a unutra  $PM_1$ ,  $PM_{2,5}$  i  $PM_{10}$  čestice pomoću GRIMM monitora. Oba su instrumenta kalibrirana gravimetrijskom metodom pomoću filtara. Kolmogorov-Smirnovljev test pokazao je normalnu raspodjelu vanjskih mjerenja. Srednje razine unutrašnjih čestica  $PM_1$ ,  $PM_{2,5}$ ,  $PM_{10}$ , odnosno vanjskih  $PM_{2,5}$  čestica, za sve su škole iznosile  $17,6 \mu\text{g m}^{-3}$ ,  $46,6 \mu\text{g m}^{-3}$ ,  $400,9 \mu\text{g m}^{-3}$ , odnosno  $36,9 \mu\text{g m}^{-3}$ . Najviše razine unutrašnjih i vanjskih  $PM_{2,5}$  čestica zabilježene u školi Shahed za dječake ( $69,1 \mu\text{g m}^{-3}$  i  $115,8 \mu\text{g m}^{-3}$ ), a najniže u školi Kazemi ( $29,1 \mu\text{g m}^{-3}$  i  $15,5 \mu\text{g m}^{-3}$ ). Bez obzira na to jesu li škole bile smještene na glavnim ili sporednim ulicama, povezanost između razina unutrašnjih sitnih čestica ( $PM_{2,5}$  i  $PM_1$ ) i razina  $PM_{2,5}$  vanjskih čestica bila je snažnija nego između razina  $PM_{10}$  čestica izmjerenih unutra i  $PM_{2,5}$  čestica izmjerenih vani. Srednje razine  $PM_{2,5}$  i  $PM_{10}$  čestica u učionicama te  $PM_{2,5}$  čestica u dvorištima škola bile su više od standarda, a razine sitnih čestica u zatvorenom i na otvorenom dobro su kolerirale.

**KLJUČNE RIJEČI:** unutrašnje čestice, vanjske čestice

## CORRESPONDING AUTHOR:

Mahmoud Mohammadyan  
Khazar Abad Road, Mazandarn University Campus  
Sari, Iran  
E-mail: mohammadyan@yahoo.com